

## Validation of Anti-GAGA4 IgM Antibodies for Predicting the Development of Relapsing Remitting Multiple Sclerosis after the First Neurological Event.

Mark S. Freedman<sup>a,\*</sup>, Jessica Laks<sup>a</sup>, Nir Dotan<sup>b</sup>, Rom T. Altstock<sup>b</sup>, Avinoam Dukler<sup>b</sup>, , and Christian Sindic<sup>c</sup>

<sup>a</sup>University of Ottawa, The Ottawa Hospital-General Campus, 501 Smyth Road, Ottawa, Ontario Canada K1H 8L6,

<sup>b</sup>Glycominds Ltd, 1 Yodfat St, Alon Bldg, Lod 71291, Israel, <sup>c</sup>Cliniques universitaires Saint-Luc, UCL, 1200 Brussels, Belgium.

**Background:** There is an unmet need to develop specific serum based biomarkers for the diagnosis and prognosis of Relapsing Remitting MS (RRMS). We have reported that elevated levels of serum anti-Glc(alpha1,4)Glc(alpha) (GAGA4) IgM antibodies (Ab) exist in RRMS (n=44) patients in comparison to patients with other neurological diseases (OND, n=44) enabling to discern which post-CIS patients convert to RRMS vs. OND. To validate our findings we have further investigate an additional and larger cohort of patients. **Aim:** To validate the predictive value of IgM Ab against GAGA4, for identifying of post CIS patients that will evolve to RRMS in a larger cohort. **Methods:** Retrospective analysis of 121 frozen sera taken from patients presenting for diagnostic work-up after a first acute neurological event consistent with demyelination (CIS) and were followed for a minimum of 4 years to confirm RRMS diagnosis. The "other neurological disease" (OND) control group comprised of 85 patients, 32 with other inflammatory disorders (OIND), and 53 with non-inflammatory neurological disease (ONIND). Sera were diluted 1:1200 and levels of anti GAGA4 IgM Ab measured by immunoassay normalized by dividing to square root of IgM levels (mg IgM/ml serum). **Results:** Significantly higher levels of anti-GAGA4 IgM/Total IgM ( $p=0.04$ , T test ) Ab were observed in CIS patients who converted to RRMS as opposed to OND. Using a cut-off of 42 (anti GAGA4 IgM EU/(mg IgM /ml serum)<sup>0.5</sup>), we have found that 34/121 (28%) converting CIS patients were positive, whereas 77/85 (91%) OND patients were negative, corresponding to a sensitivity of 28% (CI 95% 20.6-34.3%), a specificity of 91% (CI 95% 82.3-95.8%), PPV of 86.5%, and NPV of 36.4%. **Conclusion:** A higher level of Anti-GAGA4 IgM Ab was further validated as a predictor for the development of RRMS in CIS patients.

Key words: anti-glycan; antibody; biomarker, prediction

---